



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

ideal of the corn-breeder should then be continuous hybridization between biotypes, rather than the isolation of pure strains.—R. R. GATES.

**Isolation and mutation.**—While the final adjudication of the claims of the various theories of evolution must be made on an experimental basis, such data must be in harmony with the facts of plant and animal distribution, as is pointed out in a suggestive paper by LEAVITT.<sup>24</sup> It is of much interest to observe that zoologists, as a rule, have been less inclined to believe in mutation than have botanists. This is in part due, LEAVITT thinks, to a less perfect grasp of the theory by some of the zoologists, but in part due also to the fact that most students of animal distribution believe that isolation of closely related species is a most important principle in evolution. The author shows that there are innumerable cases of overlap in closely related plants of all groups, most notable, perhaps, in the widely varying thallophytes and bryophytes, but abundant in the seed plants. There is plenty of evidence that new species may have originated from the old without geographic isolation, although cases suggesting the latter are not wanting. Therefore, it is concluded, many facts of plant distribution favor the mutation theory, though they do not show that this is the only valid theory of evolution.—H. C. COWLES.

**Osmotic properties of root hairs.**—HILL<sup>25</sup> has investigated the osmotic properties of the root hairs of *Glyceria maritima*, *Suaeda maritima*, and *Salicornia herbacea*, which grow in a salt marsh subject to great changes in the osmotic pressure of its soil water, due to periodic flooding by the tides and to occasional drenching rains. He finds that the hairs show marked and rather rapid variation in osmotic pressure corresponding in variation to the osmotic pressure of the soil water. This variation is not due to the entrance of the abundant chlorids of the soil water, for in no case could he find chlorids in the root hairs, although they could be found in traces in the upper portions of the seedlings. The high osmotic pressure of the soil water seems to act through the irritability of the protoplasm, causing a dissociation of the compounds of the cells. He thinks OSTERHOUT is wrong in concluding that osmosis is not an important process in plant nutrition, and points out the fact that all the data of this investigator can be explained by the fact that plants can modify their osmotic properties readily in response to and in protection against rapidly varying external osmotic pressures.—WILLIAM CROCKER.

**Statolith theory.**—BUDER<sup>26</sup> comes to the support of the statolith theory with a set of well-chosen and critical experiments that seem to justify his conclusions,

---

<sup>24</sup> LEAVITT, R. S., The geographic distribution of nearly related species. Amer. Nat. 41:207-240. 1907.

<sup>25</sup> HILL, F. G., Observations on the osmotic properties of the root hairs of certain salt marsh plants. New Phytologist 7:133-142. 1908.

<sup>26</sup> BUDER, JOHANNES, Untersuchungen zur Statolithenhypothese. Festschrift zur Feier des 25-jährigen Bestehens der Deutsch. Bot. Gesells. Ber. 26:162-193. 1908.